— ABLYBIO, Help Your Research

TLR7 Rabbit pAb

货号: AYP14268

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	**WB:** Mus musculus
应用	WB ICC/IF
推荐浓度	**WB:** 1:500 - 1:2000 **IF/ICC:** 1:50 - 1:200
理论分子量	120kDa
实测分子量	74(cleaved)kD，100KDa-140KD
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver
细胞定位	Cytoplasmic vesicle,Endoplasmic reticulum membrane,Endosome,Lysosome,Single-pass type I membrane protein,phagosome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 891-1049 of human TLR7 (NP_057646.1).

靶点信息

研究背景	The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This gene is predominantly expressed in lung, placenta, and spleen, and lies in close proximity to another family member, TLR8, on chromosome X.
基因ID	51284
基因名	TLR7
Swiss	Q9NYK1 (https://www.uniprot.org/uniprotkb/Q9NYK1/entry)
别名	TLR7,TLR7-like,TLR7 Rabbit pAb

产品验证



Western blot analysis of TLR7 expressed in Mouse bring， Mouse lung using TLR7 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)