

— ABLYBIO, Help Your Research



ACVR1C Rabbit pAb

货号: **AYP14280**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:100 - 1:200
理论分子量	37kDa/46kDa/49kDa/54kDa
实测分子量	55KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,BxPC-3,Mouse brain,Rat brain
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 22-113 of human ACVR1C (NP_660302.2).
------	---

靶点信息

研究背景	ACVR1C is a type I receptor for the TGFB (see MIM 190180) family of signaling molecules. Upon ligand binding, type I receptors phosphorylate cytoplasmic SMAD transcription factors, which then translocate to the nucleus and interact directly with DNA or in complex with other transcription factors (Bondestam et al., 2001 [PubMed 12063393]).[supplied by OMIM, Mar 2008]
基因ID	130399
基因名	ACVR1C
Swiss	Q8NER5 (https://www.uniprot.org/uniprotkb/Q8NER5/entry)
别名	ACVR1C,ACVRLK7,ALK7,ACVR1C Rabbit pAb,Activin receptor type IC,Activin receptor-like kinase 7

产品验证



Western blot analysis of ACVR1C expressed in Mouse brain,Rat brain,HT-29 using ACVR1C Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)