

— ABLYBIO, Help Your Research



RAR α Rabbit pAb

货号: **AYP14292**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus Co-IP: Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	39kDa/50kDa
实测分子量	60KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,K-562,Raji,THP-1,SH-SY5Y,Mouse testis,Mouse brain,Rat brain
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-457 of human RAR α (NP_001019980.1).
------	---

靶点信息

研究背景	This gene represents a nuclear retinoic acid receptor. The encoded protein, retinoic acid receptor alpha, regulates transcription in a ligand-dependent manner. This gene has been implicated in regulation of development, differentiation, apoptosis, granulopoiesis, and transcription of clock genes. Translocations between this locus and several other loci have been associated with acute promyelocytic leukemia. Alternatively spliced transcript variants have been found for this locus.
基因ID	5914
基因名	RARA
Swiss	P10276 (https://www.uniprot.org/uniprotkb/P10276/entry)
别名	NR1B1,RAR,RARA,RAR α Rabbit pAb,Nuclear receptor subfamily 1 group B member 1

产品验证



Western blot analysis of RAR α expressed in Mouse testis,Rat brain,MCF7 using RAR α Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)