

ABCA7 Rabbit pAb

货号: **AYP14417**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	234kDa
实测分子量	260KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat brain
细胞定位	cell surface,endoplasmic reticulum,glial cell projection,Golgi apparatus,phagocytic cup,plasma membrane ,ruffle membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 2010-2146 of human ABCA7 (NP_061985.2).
序列	QHLKGRFAAGHTLTLRVPAARSQPAAAFVAAEFPGAELREAHGGRLRFQLPPGGRCALARVFGELAVHGAEHGVEDFSVS QTMLEEVFLYFSKDQKGDEDEEQKEAGVGVDPA PGLQHPKRVSQFLDDPSTAETVL

靶点信息

研究背景	The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the ABC1 subfamily. Members of the ABC1 subfamily comprise the only major ABC subfamily found exclusively in multicellular eukaryotes. This full transporter has been detected predominantly in myelo-lymphatic tissues with the highest expression in peripheral leukocytes, thymus, spleen, and bone marrow. The function of this protein is not yet known; however, the expression pattern suggests a role in lipid homeostasis in cells of the immune system.
基因ID	10347
基因名	ABCA7
Swiss	Q8IZY2
别名	ABCA7;ABCA-SSN;ABCX;AD9

产品验证



Western blot analysis of ABCA7 expressed in Rat brain using ABCA7 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn