

— ABLYBIO, Help Your Research



# NCEH1 Rabbit pAb

货号: **AYP14433**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	
实测分子量	46kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	PC-3,BxPC-3
细胞定位	plasma membrane
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 90-440 of human NCEH 1 (NP_065843.3).
------	---

## 靶点信息

研究背景	Hydrolyzes 2-acetyl monoalkylglycerol ether, the penultimate precursor of the pathway for de novo synthesis of platelet-activating factor. May be responsible for cholesterol ester hydrolysis in macrophages (By similarity). Also involved in organ detoxification by hydrolyzing exogenous organophosphorus compounds (By similarity). May contribute to cancer pathogenesis by promoting tumor cell migration.
基因ID	57552
基因名	NCEH1
Swiss	Q6PIU2 ( <a href="https://www.uniprot.org/uniprotkb/Q6PIU2/entry">https://www.uniprot.org/uniprotkb/Q6PIU2/entry</a> )
别名	AADACL1,NCEH,NCEH1,NCEH1 Rabbit pAb,Acetylalkylglycerol acetylhydrolase,Arylacetamide deacetylase-like 1,KIAA1363

## 产品验证



Western blot analysis of NCEH1 expressed in PC-3, BxPC-3 using NCEH1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))