

# THSD4 Rabbit pAb

货号: **AYP14435**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  | <b>WB:</b> Homo sapiens  |
| 应用    | <a href="#">WB</a>   |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000  |
| 理论分子量 |  |
| 实测分子量 | 112kDa   |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | SH-SY5Y  |
| 细胞定位  | extracellular exosome  |
| 纯化    | Affinity purification  |

## 抗原信息


|      |   |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 500-700 of human THSD4 (NP_079093.2).   |
| 序列   | AEGPTNEILDVYMIHQPNPGVHYEYVIMGTNAISPVPPHRRPGEFNGMQMVTEGRSQEEGEQKGRNEEKEDLRGEAPE<br>MFTSESAQTFPVRHPDRFSPHRPDNLVPPAPQPPRRSRDHNWKQLGTTECSTTCGKGSQYPIFRCVHRSTHEEAPESYCD<br>SSMKPTPEEEPCNIFPCPAFWDIGEWSECSKTCGLGMQHRQ |

## 靶点信息

|      |   |
|------|---|
| 研究背景 | Promotes FBN1 matrix assembly. Attenuates TGFβ signaling, possibly by accelerating the sequestration of large latent complexes of TGFβ or active TGFβ by FBN1 microfibril assembly, thereby negatively regulating the expression of TGFβ regulatory targets, such as POSTN (By similarity). |
| 基因ID | 79875   |

|       |  |
|-------|--|
| 基因名   | THSD4                                      |
| Swiss | Q6ZMP0                                     |
| 别名    | ADAMTSL-6;ADAMTSL6;FVSY9334;PRO34005;THSD4 |

## 产品验证



Western blot analysis of THSD4 expressed in SH-SY5Y using THSD4 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)