

ENAH Rabbit pAb

货号: **AYP14437**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens IP: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,C6,rat brain
细胞定位	cytoskeleton,cytosol,filopodium,focal adhesion,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 412-591 of human ENAH (NP_001008493.1).
序列	FPSGGNAIGVNSASSKTDTRGRNGPLPLGGSGLMEEEMSALLARRRRRIAIEKGSTIETEYKEDKGEDSEPVTSKASSTSTPEP TRKPWERTNTMNGSKSPVISRRDSPRKNQIVFDNRSYDSLHRPKSTPLSQPSANGVQTEGLDYDRLKQDILDEMRELTK LKEELIDAIRQELSKSNTA

靶点信息

研究背景	This gene encodes a member of the enabled/ vasodilator-stimulated phosphoprotein. Members of this gene family are involved in actin-based motility. This protein is involved in regulating the assembly of actin filaments and modulates cell adhesion and motility. Alternate splice variants of this gene have been correlated with tumor invasiveness in certain tissues and these variants may serve as prognostic markers. A pseudogene of this gene is found on chromosome 3. [provided by RefSeq, Sep 2016]
------	--

基因ID	55740
基因名	ENAH
Swiss	Q8N8S7
别名	ENA;MENA;NDPP1;ENAH

产品验证



Western blot analysis of ENAH expressed in U-251MG,C6, rat brain using ENAH Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn