

# pan-Symmetric-Di-Methyl Arginine Rabbit pAb

货号: **AYP14449**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Mus musculus
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:100 - 1:500
理论分子量	
实测分子量	22-70KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,MCF7,NIH/3T3,C6,Mouse liver,Rat liver
细胞定位	
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence containing succinylated K.
序列	

## 靶点信息

研究背景	The methylation of arginine residues in proteins is a post-translational modification that contributes to a wide range of biological processes. Many cytokines involved in T cell development and activation utilize the common cytokine receptor $\gamma$ -chain ( $\gamma$ c) and the kinase JAK3 for signal transduction.
基因ID	
基因名	
Swiss	

别名	
----	--

## 产品验证



Western blot analysis of pan-Symmetric-Di-Methyl Arginine expressed in Mouse liver,Rat liver,H eLa using pan-Symmetric-Di-Methyl Arginine Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)