

H2-Aa Rabbit pAb

货号: **AYP14452**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	28kDa
实测分子量	28KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung,Mouse kidney
细胞定位	external side of plasma membrane,lysosome,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 27-208 of mouse H2-Aa (NP_034508.2).
------	--

靶点信息

研究背景	Enables peptide antigen binding activity. Involved in response to interferon-gamma. Acts upstream of or within antigen processing and presentation of exogenous peptide antigen via MHC class II and positive regulation of T cell differentiation. Located in external side of plasma membrane and lysosome. Part of MHC class II protein complex. Is expressed in central nervous system; retina; and thymus primordium. Human ortholog(s) of this gene implicated in several diseases, including autoimmune disease (multiple); gastrointestinal system cancer (multiple); glomerulonephritis (multiple); hematologic cancer (multiple); and systemic sclerosis (multiple). Orthologous to several human genes including HLA-DQA1 (major histocompatibility complex, class II, DQ alpha 1).
基因ID	14960

基因名	H2-Aa
Swiss	P14434
别名	Ia1;H2Aa;Ia-1;H-2Aa;Aalpha;IAalpha;I-Aalpha;H2-Aa

产品验证



Western blot analysis of H2-Aa expressed in Mouse lung, Mouse kidney using H2-Aa Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn