

— ABLYBIO, Help Your Research



TPX2 Rabbit pAb

货号: **AYP14471**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	IF: Homo sapiens
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	85kDa
实测分子量	105KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549
细胞定位	Cytoplasm,Nucleus,cytoskeleton,spindle,spindle pole
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human TPX2 (N P_036244.2).
------	---

靶点信息

研究背景	Spindle assembly factor required for normal assembly of mitotic spindles. Required for normal assembly of microtubules during apoptosis. Required for chromatin and/or kinetochore dependent microtubule nucleation. Mediates AURKA localization to spindle microtubules. Activates AURKA by promoting its autophosphorylation at 'Thr-288' and protects this residue against dephosphorylation. TPX2 is inactivated upon binding to importin-alpha. At the onset of mitosis, GOLGA2 interacts with importin-alpha, liberating TPX2 from importin-alpha, allowing TPX2 to activate AURKA kinase and stimulate local microtubule nucleation.
基因ID	22974
基因名	TPX2
Swiss	Q9ULW0 (https://www.uniprot.org/uniprotkb/Q9ULW0/entry)
别名	DIL2,p100,DIL-2,HCTP4,FLS353,HCA519,REPP86,C20orf1,C20orf2,GD:C20orf1,TPX2,TPX2 Rabbit pAb,Differentially expressed in cancerous and non-cancerous lung cells 2,Hepatocellular carcinoma-associated antigen 519,Hepatocellular carcinoma-associated antigen 90,Protein fls353

产品验证



Western blot analysis of TPX2 expressed in A-549 using TPX2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)