

# MAN2A2 Rabbit pAb

货号: **AYP14509**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	<b>WB:</b> Homo sapiens
应用	<a href="#">WB</a>
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	
实测分子量	128KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG
细胞定位	
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human MAN2A2 (NP_006113.2).
序列	VPPEPRPSFFSISPQDCQFALGGRGQKPELQMLTVSEELPFDNVDGGVWRQGFDISYDPHDWDAEDLQVFVPHSHND PGWIKTFDKYYTEQTQHILNSMV

## 靶点信息

研究背景	Catalyzes the first committed step in the biosynthesis of complex N-glycans. It controls conversion of high mannose to complex N-glycans; the final hydrolytic step in the N-glycan maturation pathway.
基因ID	4122
基因名	MAN2A2

Swiss	P49641
别名	MANA2X;MAN2A2

## 产品验证



Western blot analysis of MAN2A2 expressed in U-87MG using MAN2A2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)