

— ABLYBIO, Help Your Research



PDIA4 Rabbit pAb

货号: **AYP14949**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	72kDa
实测分子量	72KDa
形式	Liquid
保存条件	Store at -20°C Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,MCF7,Mouse liver,Rat lung,Rat liver
细胞定位	Endoplasmic reticulum lumen,Melanosome
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 200-300 of human PDIA4 (NP_00490 2.1).
------	---

靶点信息

研究背景	This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, three catalytically active thioredoxin (TRX) domains, two TRX-like domains and a C-terminal ER-retention sequence. This protein, when bound to cyclophilin B, enhances the rate of immunoglobulin G intermolecular disulfide bonding and antibody assembly.
基因ID	9601
基因名	PDIA4
Swiss	P13667 (https://www.uniprot.org/uniprotkb/P13667/entry)
别名	PDIA4,ERP70,ERP72,ERp-72,PDIA4 Rabbit pAb,Endoplasmic reticulum resident protein 70,Endoplasmic reticulum resident protein 72

产品验证



Western blot analysis of PDIA4 expressed in Mouse liver, Rat lung, HepG2 using PDIA4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)