

Phospho-TrkA-Y674/675 + TrkB-Y706/707 Rabbit pAb

货号: AYP15048

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:100 - 1:500
理论分子量	35kDa/53-60kDa/77kDa/83-93kDa
实测分子量	140KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	axon,dendrite,early endosome,late endosome,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	A synthetic phosphorylated peptide around Y674 & Y675 of human TrkANTRK1 (NP_001012331.1).
序列	TDYYRV

靶点信息

研究背景	This gene encodes a member of the neurotrophic tyrosine kinase receptor (NTRK) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MA PK pathway. The presence of this kinase leads to cell differentiation and may play a role in specifying sensory neuron subtypes. Mutations in this gene have been associated with congenital insensitivity to pain, anhidrosis, self-mutilating behavior, mental retardation and cancer. Alternate transcriptional splice variants of this gene have been found, but only three have been characterized to date. [provided by RefSeq, Jul 2008]
------	--

基因ID	4914,4915
基因名	NTRK1,NTRK2
Swiss	P04629,Q16620
别名	NTRK1/NTRK2

产品验证



Western blot analysis of Phospho-TrkA-Y674/675 + TrkB-Y706/707 expressed in HeLa using Phospho-TrkA-Y674/675 + TrkB-Y706/707 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn