

— ABLYBIO, Help Your Research



Phospho-HDAC4-S632 Rabbit pAb

货号: **AYP15082**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	106kDa/119kDa
实测分子量	140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A phospho specific peptide corresponding to residues surrounding S632 of human HDAC4
------	--

靶点信息

研究背景	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3.
基因ID	9759
基因名	HDAC4
Swiss	P56524 (https://www.uniprot.org/uniprotkb/P56524/entry)
别名	HDAC4,AHO3,BDMR,HA6116,HD4,HDAC-4,HDAC-A,HDACA,Phospho-HDAC4-S632 Rabbit pAb,KIAA0288

产品验证



Western blot analysis of Phospho-HDAC4-S632 expressed in 293 using Phospho-HDAC4-S632 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)