— ABLYBIO, Help Your Research

Phospho-PRKCQ-S676 Rabbit pAb

货号: AYP15107

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	**WB:** 1:500 - 1:1000 **IHC:** 1:50 - 1:100 **IF/ICC:** 1:100 - 1:200
理论分子量	67kDa/74kDa/81kDa
实测分子量	78KDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat
细胞定位	Cell membrane,Cytoplasm,Peripheral membrane protein
纯化	Affinity purification


抗原信息

抗原信息	A phospho specific peptide corresponding to residues surrounding S676 of human PRKCQ

靶点信息

研究背景	Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role. The protein encoded by this gene is one of the PKC family members. It is a calcium-independent and phospholipid-dependent protein kinase. This kinase is important for T-cell activation. It is required for the activation of the transcription factors NF-kappaB and AP-1, and may link the T cell receptor (TCR) signaling complex to the activation of the transcription factors.
基因ID	5588
基因名	PRKCQ
Swiss	Q04759 (https://www.uniprot.org/uniprotkb/Q04759/entry)
别名	PRKCQ,PRKCT,nPKC-theta,Phospho-PRKCQ-S676 Rabbit pAb

产品验证



Western blot analysis of Phospho-PRKCQ-S676 expressed in Jurkat using Phospho-PRKCQ-S676 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)