

NECAB1 Rabbit pAb

货号: **AYP15113**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	11kDa/40kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-351 of human NECAB 1 (NP_071746.1).
序列	MEDSQETSPSSNNSSEELSSALHLSKGMISIFLDILRRADKNDGKLSFEFEKAYFADGVLSGEELHELFTIDHTNTNLD TEELCEYFSQHLGEYENVLAALEDLNLKAMGKTKKDYQEASNLQFVTRFLLKETLNQLQSLQNSLECAMETTEEQTRQ ERQGPAKPEVLSIQWPGKRSSRRVQRHNSFSPNSPQFNVSQGLLEEDNQWMTQINRLQKLDRLEKKDLKLEPPEEEIIE GNTKSHIMLVQRQMSVIEEDLEEFQLALKHYVESASSQSGCLRISIQKLSNESRYMIYEFWENSSVWNSHLQTNYSKTFQR SNVDFLETPELTSTMLVPASWWILNN

靶点信息

研究背景	Enables identical protein binding activity. Predicted to be involved in regulation of amyloid precursor protein biosynthetic process. Predicted to act upstream of or within blastocyst hatching. Located in cytosol and nucleoplasm.
------	---

基因ID	64168
基因名	NECAB1
Swiss	Q8N987
别名	NECAB1;EFCBP1;STIP-1

产品验证



Western blot analysis of NECAB1 expressed in LO2 using NECAB1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn