

— ABLYBIO, Help Your Research



# SNAPC2 Rabbit pAb

货号: **AYP15122**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	35kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 20-100 of human SNAP C2 (NP_003074.1).
------	--

## 靶点信息

研究背景	This gene encodes a subunit of the snRNA-activating protein complex which is associated with the TATA box-binding protein. The encoded protein is necessary for RNA polymerase II and III dependent small-nuclear RNA gene transcription. Alternative splicing results in multiple transcript variants.
基因ID	6618
基因名	SNAPC2
Swiss	Q13487 ( <a href="https://www.uniprot.org/uniprotkb/Q13487/entry">https://www.uniprot.org/uniprotkb/Q13487/entry</a> )
别名	SNAPC2,PTFDELTA,SNAP45,SNAPC2 Rabbit pAb,Proximal sequence element-binding transcription factor subunit delta,Small nuclear RNA-activating complex polypeptide 2,snRNA-activating protein complex 45 kDa subunit

## 产品验证



Western blot analysis of SNAPC2 expressed in CACO2, mouse brain using SNAPC2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))