

— ABLYBIO, Help Your Research



MED25 Rabbit pAb

货号: **AYP15124**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	55kDa/76kDa/78kDa/82kDa/84kDa
实测分子量	78kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,A-549,mouse brain, rat kidney
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 30-200 of human MED25 (NP_112235.2).
------	--

靶点信息

研究背景	This gene encodes a component of the transcriptional coactivator complex termed the Mediator complex. This complex is required for transcription of most RNA polymerase II-dependent genes. The encoded protein plays a role in chromatin modification and in preinitiation complex assembly. Mutations in this gene are associated with Charcot-Marie-Tooth disease type 2B2.
基因ID	81857
基因名	MED25
Swiss	Q71SY5 (https://www.uniprot.org/uniprotkb/Q71SY5/entry)
别名	MED25,ACID1,ARC92,BVSY5,CMT2B2,P78,PTOV2,TCBAP0758,MED25 Rabbit pAb,Activator interaction domain-containing protein 1,Activator-recruited cofactor 92 kDa component,Mediator complex subunit 25

产品验证



Western blot analysis of MED25 expressed in mouse brain, rat kidney, K-562 using MED25 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)