

# MED25 Rabbit pAb

货号: AYP15124

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	55kDa/76kDa/78kDa/82kDa/84kDa
实测分子量	78kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,A-549,mouse brain,rat kidney
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 30-200 of human MED25 (NP_112235.2).
序列	PYFEGLRKHILLPAIEYFNGGPPAETDFGGDYGGTQYSLVVFNTVDCAPESYVQCHAPTSSAYEFVTWLDGIKFMGGGGE SCSLIAEGLSTALQLFDDFKMREQIQTHRVCLLICNSPPYLLPAVESTTYSGCTTENLVQQIGERGIHFSIVSPRKLPAALRL FEKAAP

## 靶点信息

研究背景	This gene encodes a component of the transcriptional coactivator complex termed the Mediator complex. This complex is required for transcription of most RNA polymerase II-dependent genes. The encoded protein plays a role in chromatin modification and in preinitiation complex assembly. Mutations in this gene are associated with Charcot-Marie-Tooth disease type 2B2.
------	--

基因ID	81857
基因名	MED25
Swiss	Q71SY5
别名	MED25;ACID1;ARC92;BVSYS;CMT2B2;P78;PTOV2;TCBAP0758

## 产品验证



Western blot analysis of MED25 expressed in mouse brain, rat kidney, K-562 using MED25 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)