

MNAT1 Rabbit pAb

货号: **AYP15138**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:50 - 1:200
理论分子量	31kDa/35kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	PC-3,22Rv1,DU 145,293T,Mouse testis,Mouse liver
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-309 of human MNAT1 (NP_002422.1).
序列	MDDQGCPRCKTTKYRNP SLKLMVNVCGHTLCESCVDLLFVRGAGNCPECGTPLRKS NFRVQLFEDPTVDKEVEIRKVKLKI YNKREEDFPSLREYND FLEEV EIVFNLTNNVDLNTKKKMEIYQKENKDVIQKNK LKLTREQEELEEALEVERQENEQRRLF IQKEEQ LQQLKRKNKQAF LDELESSDLPVALLLAQHKDRSTQLEMQLKPKPKVPTVFSTGKIMGQHISLAPIHKLEEA LYE YQPLQIETYGPHVPELEMLGRLGYLNHVRAASPQDLAGGYTSSLACHRALQDAFSGLFWQPS

靶点信息

研究背景	The protein encoded by this gene, along with cyclin H and CDK7, forms the CDK-activating kinase (CAK) enzymatic complex. This complex activates several cyclin-associated kinases and can also associate with TFIID to activate transcription by RNA polymerase II. Two transcript variants encoding different isoforms have been found for this gene.
------	--

基因ID	4331
基因名	MNAT1
Swiss	P51948
别名	MNAT1;CAP35;MAT1;RNF66;TFB3

产品验证



Western blot analysis of MNAT1 expressed in Mouse testis, PC-3, 22Rv1 using MNAT1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn