

# BCAS2 Rabbit pAb

货号: **AYP15174**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	26kDa
实测分子量	25kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Jurkat,MCF-7,HeLa,SKOV3,Mouse testis,Mouse spleen,Mouse lung,Rat lung
细胞定位	Nucleus,nucleolus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-225 of human BCAS2 (NP_005863.1).
序列	MAGTGLVAGEVVVDALPYFDQGYEAPGVREAAAALVEEETRYYRPTKNYLSYLTAPDYSAFETDIMRNEFERLAARQPIELL SMKRYELPAPSSGQKNDITAWQECVNNNSMAQLEHQAVRIENLELMSQHGCNAWKVYNNENLVHMIEHAQKELQKLRKHIQ DLNWQRKNMQLTAGSKLREMESNWVSLVSKNYEIERIVQLENEIYQIKQQHGEANKENIRQDF

## 靶点信息

研究背景	Required for pre-mRNA splicing as component of the activated spliceosome. Component of the PRP19-CD C5L complex that forms an integral part of the spliceosome and is required for activating pre-mRNA splicing. May have a scaffolding role in the spliceosome assembly as it contacts all other components of the core complex. The PRP19-CDC5L complex may also play a role in the response to DNA damage (DDR).
------	---

基因ID	10286
基因名	BCAS2
Swiss	O75934
别名	BCAS2;DAM1;SPF27;Snt309

## 产品验证



Western blot analysis of BCAS2 expressed in Mouse testis,Rat lung,293T using BCAS2 Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)