

— ABLYBIO, Help Your Research



UGT1A4 Rabbit pAb

货号: **AYP15198**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	49kDa/60kDa
实测分子量	69kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,A-549,HT-29,HepG2
细胞定位	Endoplasmic reticulum membrane,Microsome,Single-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 29-150 of human UGT1A4 (NP_009051.1).
------	---------------------------------------------------------------------------------------------------------------------

靶点信息

研究背景	This gene encodes a UDP-glucuronosyltransferase, an enzyme of the glucuronidation pathway that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into water-soluble, excretable metabolites. This gene is part of a complex locus that encodes several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons followed by four common exons. Four of the alternate first exons are considered pseudogenes. Each of the remaining nine 5' exons may be spliced to the four common exons, resulting in nine proteins with different N-termini and identical C-termini. Each first exon encodes the substrate binding site, and is regulated by its own promoter. This enzyme has some glucuronidase activity towards bilirubin, although it is more active on amines, steroids, and sapogenins.
基因ID	54657
基因名	UGT1A4
Swiss	P22310 (https://www.uniprot.org/uniprotkb/P22310/entry)
别名	UGT1A4,HUG-BR2,UDPGT,UDPGT 1-4,UGT-1D,UGT1-04,UGT1.4,UGT1A4S,UGT1D,UDPGT1-4,UGT1A4 Rabbit pAb,Bilirubin-specific UDPGT isozyme 2,UDP-glucuronosyltransferase 1-4,UDP-glucuronosyltransferase 1-D,GNT1,UGT1

产品验证



Western blot analysis of UGT1A4 expressed in MCF7, A-549, HT-29 using UGT1A4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)