

— ABLYBIO, Help Your Research



# WDR48 Rabbit pAb

货号: **AYP15255**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:100
理论分子量	13kDa/45kDa/66kDa/75kDa/76kDa
实测分子量	76kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,A-431,SGC-7901,LO2,A-549,HT-1080,Mouse brain,Rat testis
细胞定位	Cytoplasm,Lysosome,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 450-677 of human WDR 48 (NP_065890.1).
------	--

## 靶点信息

研究背景	The protein encoded by this gene has been shown to interact with ubiquitin specific peptidase 1 (USP1), activating the deubiquitinating activity of USP1 and allowing it to remove the ubiquitin moiety from mono ubiquitinated FANCD2. FANCD2 is ubiquitinated in response to DNA damage.
基因ID	57599
基因名	WDR48
Swiss	Q8TAF3 ( <a href="https://www.uniprot.org/uniprotkb/Q8TAF3/entry">https://www.uniprot.org/uniprotkb/Q8TAF3/entry</a> )
别名	WDR48,P80,SPG60,UAF1,WDR48 Rabbit pAb,USP1-associated factor 1,WD repeat endosomal protein,KIAA 1449

## 产品验证



Western blot analysis of WDR48 expressed in Mouse brain,Rat testis,U-251MG using WDR48 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))