

SS18 Rabbit pAb

货号: **AYP15267**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	42kDa/45kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,Jurkat
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-240 of human SS18 (NP_001007560.1).
序列	MSVAF AAPRQRGKGEITPA AIQKMLDDNNHLIQCIMDSQNKGKTSECSQYQQLHTNLVYLATIADSNQNMQSLLPAPPT QNMPMGPGGMNQSGPPPPRSHNMPSDGMVGGGPPAPHMQNQMNQMPGPNHMPMQGPGPNQLNMTNSSMNMP SSSHGSMGGYNHVPSSQSMPVQNMQMTMSQGQPMGNYGPRPNMSMQPNQGPMHQPPSQYQNPQGGGQHYQ GQPPMGMMGQVNVQ

靶点信息

研究背景	Appears to function synergistically with RBM14 as a transcriptional coactivator. Isoform 1 and isoform 2 function in nuclear receptor coactivation. Isoform 1 and isoform 2 function in general transcriptional coactivation. Component of SWI/SNF chromatin remodeling subcomplex GBAF that carries out key enzymatic activities, changing chromatin structure by altering DNA-histone contacts within a nucleosome in an ATP-dependent manner.
------	--

基因ID	6760
基因名	SS18
Swiss	Q15532
别名	SS18;SSXT;SYT

产品验证



Western blot analysis of SS18 expressed in MCF7, Jurkat using SS18 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn