

STX8 Rabbit pAb

货号: **AYP15337**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	26kDa
实测分子量	27kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,293T,HepG2,SW480,Mouse lung,Mouse kidney,Mouse liver
细胞定位	Membrane,Single-pass type IV membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-215 of human STX8 (NP_004844.1).
序列	MAPDPWFSTYDSTCQIAQEIAEKIQQRNQYERKGEKAPKLTVTIRALLQNLKEKIALLDLLLRAVSTHQITQLEGDRRQNL DDLVTRELLLASFKNEGAEPDLIRSSLMSEEAKRGAPNPWLFEEPEETRGLGFDEIRQQQKIIQEIQDAGLDALSSISRQ KQMGQEIGNELDEQNEIIDDLANLVENTDEKLRNETRRVNMVDRKSASCG

靶点信息

研究背景	The gene is a member of the syntaxin family. The encoded protein is involved in protein trafficking from early to late endosomes via vesicle fusion and exocytosis. A related pseudogene has been identified on chromosome 12. Alternative splicing results in multiple transcript variants.
基因ID	9482

基因名	STX8
Swiss	Q9UNK0
别名	STX8;CARB

产品验证



Western blot analysis of STX8 expressed in Mouse lung,SH-SY5Y,293T using STX8 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn