

— ABLYBIO, Help Your Research



# RORC Rabbit pAb

货号: **AYP15343**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	55kDa/58kDa
实测分子量	58KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Molt-4
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 223-312 of human RORC (NP_005051.2).
------	--

## 靶点信息

研究背景	The protein encoded by this gene is a DNA-binding transcription factor and is a member of the NR1 subfamily of nuclear hormone receptors. The specific functions of this protein are not known; however, studies of a similar gene in mice have shown that this gene may be essential for lymphoid organogenesis and may play an important regulatory role in thymopoiesis. In addition, studies in mice suggest that the protein encoded by this gene may inhibit the expression of Fas ligand and IL2. Two transcript variants encoding different isoforms have been found for this gene.
基因ID	6097
基因名	RORC
Swiss	P51449 ( <a href="https://www.uniprot.org/uniprotkb/P51449/entry">https://www.uniprot.org/uniprotkb/P51449/entry</a> )
别名	RORC,IMD42,NR1F3,RORG,RZR-GAMMA,RZRG,TOR,RORC Rabbit pAb,Nuclear receptor RZR-gamma,Nuclear receptor subfamily 1 group F member 3,RAR-related orphan receptor C,Retinoid-related orphan receptor-gamma

## 产品验证



Western blot analysis of RORC expressed in Molt-4 using RORC Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)