

— ABLYBIO, Help Your Research



IL20RA Rabbit pAb

货号: **AYP15347**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	49kDa/57kDa/62kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,U-87MG,A375,SW620,Mouse liver
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 30-250 of human IL20RA (Q9UHF4).
------	--

靶点信息

研究背景	This gene encodes a member of the type II cytokine receptor family. The encoded protein is a subunit of the receptor for interleukin 20, a cytokine that may be involved in epidermal function. The interleukin 20 receptor is a heterodimeric complex consisting of the encoded protein and interleukin 20 receptor beta. This gene and interleukin 20 receptor beta are highly expressed in skin, and are upregulated in psoriasis. Alternative splicing results in multiple transcript variants.
基因ID	53832
基因名	IL20RA
Swiss	Q9UHF4 (https://www.uniprot.org/uniprotkb/Q9UHF4/entry)
别名	IL20RA,CRF2-8,IL-20R-alpha,IL-20R1,IL-20RA,IL20RA Rabbit pAb,Cytokine receptor class-II member 8,Cytokine receptor family 2 member 8,ZcytoR7

产品验证



Western blot analysis of IL20RA expressed in Mouse liver, MCF7, U-87MG using IL20RA Rabbit pAb at 1:100. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)