

— ABLYBIO, Help Your Research



# ZC3H14 Rabbit pAb

货号: **AYP15357**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	32-34kDa/61-66kDa/80-82kDa
实测分子量	105kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,A-549,HT-1080,293T
细胞定位	Cytoplasm,Nucleus speckle
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-306 of human ZC3H14 (NP_997545.2).
------	--

## 靶点信息

研究背景	The protein encoded by this gene is a poly(A)-binding protein that can affect gene expression and poly(A) tail length. The encoded protein may influence mRNA stability, nuclear export, and translation.
基因ID	79882
基因名	ZC3H14
Swiss	Q6PJT7 ( <a href="https://www.uniprot.org/uniprotkb/Q6PJT7/entry">https://www.uniprot.org/uniprotkb/Q6PJT7/entry</a> )
别名	ZC3H14,MRT56,MSUT-2,NY-REN-37,SUT2,UKp68,ZC3H14 Rabbit pAb,Mammalian suppressor of tau pathology-2,Renal carcinoma antigen NY-REN-37

## 产品验证



Western blot analysis of ZC3H14 expressed in MCF7,A-549,HT-1080 using ZC3H14 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))