

AOAH Rabbit pAb

货号: **AYP15358**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	61kDa/65kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 306-575 of human AOAH (NP_001628.1).
序列	GFLDSTVGIKEKSIYLRRLWKRNHCHNRDYNISRNGASSRNLLKFFIESLSRNKVLDPYPAIVIAMIGNDVCSGKSDPVPAMT TPEKLYSNVMQTLKHLNSHLPNGSHVILYGLPDGTFLWDLNHNRYHPLGQLNKDMTYAQLYSFLNCLQVSPCHGWMSSN KTLRRLTSEAEQLSNTLKKIAASEKFTNFNLFYMDFAFHEIIQEWQKRGQPWQLIEPVDGFHPNEVALLLLADHFWKKV QLQWPQILGKENPFNPQIKQVFGDQGGH

靶点信息

研究背景	This locus encodes both the light and heavy subunits of acyloxyacyl hydrolase. The encoded enzyme catalyzes the hydrolysis of acyloxyacyl-linked fatty acyl chains from bacterial lipopolysaccharides, effectively detoxifying these molecules. The encoded protein may play a role in modulating host inflammatory response to gram-negative bacteria. Alternatively spliced transcript variants have been described.
------	--

基因ID	313
基因名	AOAH
Swiss	P28039
别名	AOAH

产品验证



Western blot analysis of AOAH expressed in BxPC-3 using AOAH Rabbit pAb at 1:1000. Secondary antibody y: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn