

CFHR5 Rabbit pAb

货号: **AYP15359**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 64kDa |
| 实测分子量 | 75kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HepG2,K-562,Mouse liver,Rat liver |
| 细胞定位 | Secreted |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 450-569 of human CFHR5 (NP_110414.1). |
| 序列 | GPPPSINNGDTSFPLSVYPPGSTVTYRCQSFYKLGSVTVTCRNKQWSEPPRCLDPCVVSEENMNKNNIQLKWRNDGK LYAKTGDAVEFQCKFPHKAMISSPPFRAICQEGKFEYPICE |

靶点信息

| | |
|-------|---|
| 研究背景 | This gene is a member of a small complement factor H (CFH) gene cluster on chromosome 1. Each member of this gene family contains multiple short consensus repeats (SCRs) typical of regulators of complement activation. The protein encoded by this gene has nine SCRs with the first two repeats having heparin binding properties, a region within repeats 5-7 having heparin binding and C reactive protein binding properties, and the C-terminal repeats being similar to a complement component 3 b (C3b) binding domain. This protein co-localizes with C3, binds C3b in a dose-dependent manner, and is recruited to tissues damaged by C-reactive protein. Allelic variations in this gene have been associated, but not causally linked, with two different forms of kidney disease: membranoproliferative glomerulonephritis type II (MPGNII) and hemolytic uremic syndrome (HUS). |
| 基因ID | 81494 |
| 基因名 | CFHR5 |
| Swiss | Q9BXR6 |
| 别名 | CFHR5;CFHL5;CFHR5D;FHR-5;FHR5 |

产品验证



Western blot analysis of CFHR5 expressed in Mouse liver, Rat liver, HepG2 using CFHR5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn