

— ABLYBIO, Help Your Research



# Phospho-ULK1-S467 Rabbit pAb

货号: **AYP15372**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:1000
理论分子量	112kDa
实测分子量	140KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat
细胞定位	Cytoplasm,Preautophagosomal structure,cytosol
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic phosphorylated peptide around S467 of human ULK1 (NP_003556.2).
------	---

## 靶点信息

研究背景	Serine/threonine-protein kinase involved in autophagy in response to starvation. Acts upstream of phosphatidylinositol 3-kinase PIK3C3 to regulate the formation of autophagosomes, the precursors of autophagosomes. Part of regulatory feedback loops in autophagy: acts both as a downstream effector and negative regulator of mammalian target of rapamycin complex 1 (mTORC1 via interaction with RPTOR. Activated via phosphorylation by AMPK and also acts as a regulator of AMPK by mediating phosphorylation of AMPK subunits PRKAA1, PRKAB2 and PRKAG1, leading to negatively regulate AMPK activity. May phosphorylate ATG13/KIAA0652 and RPTOR; however such data need additional evidences. Plays a role early in neuronal differentiation and is required for granule cell axon formation. May also phosphorylate SESN2 and SQSTM1 to regulate autophagy. Phosphorylates FLCN, promoting autophagy. Phosphorylates AMBRA1 in response to autophagy induction, releasing AMBRA1 from the cytoskeletal docking site to induce autophagosome nucleation.
基因ID	8408
基因名	ULK1
Swiss	O75385 ( <a href="https://www.uniprot.org/uniprotkb/O75385/entry">https://www.uniprot.org/uniprotkb/O75385/entry</a> )
别名	ULK1,ATG1,ATG1A,UNC51,Unc51.1,hATG1,Phospho-ULK1-S467 Rabbit pAb,Autophagy-related protein 1 homolog,Unc-51-like kinase 1,KIAA0722

## 产品验证



Western blot analysis of Phospho-ULK1-S467 expressed in HeLa, Jurkat using Phospho-ULK1-S467 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)