

— ABLYBIO, Help Your Research



# DOCK4 Rabbit pAb

货号: **AYP15378**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	29kDa/222kDa/225kDa/226kDa
实测分子量	235kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	DU145
细胞定位	Cytoplasm, Endomembrane system, Peripheral membrane protein, cytosol
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 150-450 of human DOCK4 (NP_055520.3).
------	---

## 靶点信息

研究背景	This gene is a member of the dedicator of cytokinesis (DOCK) family and encodes a protein with a DHR-1 (CZH-1) domain, a DHR-2 (CZH-2) domain and an SH3 domain. This membrane-associated, cytoplasmic protein functions as a guanine nucleotide exchange factor and is involved in regulation of adherens junctions between cells. Mutations in this gene have been associated with ovarian, prostate, glioma, and colorectal cancers. Alternatively spliced variants which encode different protein isoforms have been described, but only one has been fully characterized.
基因ID	9732
基因名	DOCK4
Swiss	Q8N1I0 ( <a href="https://www.uniprot.org/uniprotkb/Q8N1I0/entry">https://www.uniprot.org/uniprotkb/Q8N1I0/entry</a> )
别名	DOCK4,DOCK4 Rabbit pAb,KIAA0716

## 产品验证



Western blot analysis of DOCK4 expressed in DU145 using DOCK4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)