

EPHA3/EPHA4/EPHA5 pAb

货号: **AYP15389**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	IHC IF/ICC
推荐浓度	IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	actin cytoskeleton,cytosol,early endosome,extracellular region,nuclear membrane,nucleoplasm,plasma membrane
纯化	Affinity purification

抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 700-800 of human EPHA3/EPHA4/EPHA5 (NP_005224.2).
序列	EYMENGLDLSFLRKHDAQFTVIQLVGMRLGIASGMKYLSDMGYVHRDLAARNILINSLVCKVSDFGLSRVLEDDPEAA Y TTRGGKIPRWTSPEAIAYRK

靶点信息

研究背景	Receptor tyrosine kinase which binds promiscuously membrane-bound ephrin family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Highly promiscuous for ephrin-A ligands it binds preferentially EFNA5. Upon activation by EFNA5 regulates cell-cell adhesion, cytoskeletal organization and cell migration. Plays a role in cardiac cells migration and differentiation and regulates the formation of the atrioventricular canal and septum during development probably through activation by EFNA1. Involved in the retinotectal mapping of neurons. May also control the segregation but not the guidance of motor and sensory axons during neuromuscular circuit development.
基因ID	2042,2043,2044
基因名	EPHA3,EPHA4,EPHA5
Swiss	P29320,P54764,P54756
别名	

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn