

ABR Rabbit pAb

货号: **AYP15391**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	35kDa/92kDa/93kDa/97kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse thymus,Rat lung,Rat spleen
细胞定位	axon,cytosol,dendritic spine,glutamatergic synapse,Schaffer collateral - CA1 synapse
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 200 to the C-terminus of human ABR (NP_001243776.1).
序列	DPAAKENCMMHLLRSLPDPNLITFLFLEHLKRVAEKEPINKMSLHNLATVFGPTLLRPSEVESKAHLTSAADIWSHDVMAQ VQVLLYYLQHPPISFAELKRNTLYFSTDV

靶点信息

研究背景	This gene encodes a protein that is similar to the protein encoded by the breakpoint cluster region gene 1 located on chromosome 22. The protein encoded by this gene contains a GTPase-activating protein domain, a domain found in members of the Rho family of GTP-binding proteins. Functional studies in mice determined that this protein plays a role in vestibular morphogenesis. Alternatively spliced transcript variants have been reported for this gene.
基因ID	29

基因名	ABR
Swiss	Q12979
别名	ABR;MDB

产品验证



Western blot analysis of ABR expressed in Mouse thymus,Rat lung,Rat spleen using ABR Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn