

DIXDC1 Rabbit pAb

货号: **AYP15398**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:200 - 1:2000
理论分子量	24kDa/53kDa/77kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	MCF7, 293T
细胞定位	Cell junction, Cytoplasm, cytosol, focal adhesion
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 173-472 of human DIXDC1 (NP_219493.1).
序列	SVLQLKQELLRANMDKDELHNQNVLDLQRKLDERNRLLGEYKKELGQKDRLLQQHQAKLEEALRKLSDVSYHQVDLEREL EHKDVLLAHCМКREADEATNYNSHNSQSNGLPTAGKGATSVSNRGTSDLQLVRDALRSLRNSFSGHDPQHHTIDSLE QGISSLMERLHVMETQKKQERKVRVKSPTQVGSEYRESWPPNSKLPHSQSSPTVSSTCTKVLYFTDRSLTPFMVNIPKRL EEVTLKDFKAAIDREGNHRHYHFKALDPEFGTVKEEIFHDDDAIPGWEGKIVAWVEEDHGEN

靶点信息

研究背景	The protein encoded by this gene is a positive regulator of the Wnt signaling pathway. The encoded protein is found associated with gamma tubulin at the centrosome. Alternative splicing results in multiple transcript variants encoding different isoforms.
基因ID	85458

基因名	DIXDC1
Swiss	Q155Q3
别名	DIXDC1;CCD1;dixin

产品验证



Western blot analysis of DIXDC1 expressed in MCF7,293T using DIXDC1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn