

RELT Rabbit pAb

货号: **AYP15399**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:200 - 1:2000
理论分子量	46kDa
实测分子量	47kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,K-562,LO2,Mouse thymus,Mouse brain,Rat thymus,Rat brain
细胞定位	Cell membrane,Cytoplasm,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 26-162 of human RELT (NP_116260.2).
序列	STTLWQCPPGEEPDLDPGQGTLCRPCPPGTFSAAWGSSPCQP HARCSLWRRLEAQVGMATRD TLCGDCWPGWFGPW GVPRVPCQPCSWAPLGTHGCDEWGRRARRGVEVAAGASSGGETRQPGNGTRAGGPEETAAQ

靶点信息

研究背景	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is especially abundant in hematologic tissues. It has been shown to activate the NF-kappaB pathway and selectively bind TNF receptor-associated factor 1 (TRAF1). This receptor is capable of stimulating T-cell proliferation in the presence of CD3 signaling, which suggests its regulatory role in immune response. Two alternative spliced transcript variants of this gene encoding the same protein have been reported.
基因ID	84957

基因名	RELT
Swiss	Q969Z4
别名	RELT;TNFRSF19L;TRLT

产品验证



Western blot analysis of RELT expressed in Mouse thymus,Rat thymus,Raji using RELT Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn