

# FHOD1 Rabbit pAb

货号: **AYP15409**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  |  |
| 应用    | WB   |
| 推荐浓度  | <b>WB:</b> 1:200 - 1:2000  |
| 理论分子量 | 126kDa   |
| 实测分子量 | 145kDa   |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | HeLa   |
| 细胞定位  | Cell projection, Cytoplasm, bleb, cytoskeleton   |
| 纯化    | Affinity purification  |

## 抗原信息


|      |   |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1010-1120 of human FHOD1 (NP_037373.2). |
| 序列   | TRGRMITETEKFSGVAGEAPSNPSVPAVSSGPGRGDADSHASKLLTSRPEDTTHNRRSRGMVQSSSPIMPTVGPSTASPEEPPGSSLPDTSDEIMDLLVQSVTKSSP           |

## 靶点信息

|      |   |
|------|---|
| 研究背景 | This gene encodes a protein which is a member of the formin/diaphanous family of proteins. The gene is ubiquitously expressed but is found in abundance in the spleen. The encoded protein has sequence homology to diaphanous and formin proteins within the Formin Homology (FH)1 and FH2 domains. It also contains a coiled-coil domain, a collagen-like domain, two nuclear localization signals, and several potential PKC and PKA phosphorylation sites. It is a predominantly cytoplasmic protein and is expressed in a variety of human cell lines. Alternative splicing results in multiple transcript variants. |
|------|---|

|       |            |
|-------|------------|
| 基因ID  | 29109      |
| 基因名   | FHOD1      |
| Swiss | Q9Y613     |
| 别名    | FHOD1;FHOS |

## 产品验证



Western blot analysis of FHOD1 expressed in HeLa using FHOD1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)