

FLII Rabbit pAb

货号: **AYP15427**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:200 - 1:2000 IHC: 1:50 - 1:200
理论分子量	138kDa/143kDa/144kDa
实测分子量	160kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,293T,HeLa,Jurkat
细胞定位	Cell junction,Cytoplasm,Nucleus,centrosome,cytoskeleton,focal adhesion,microtubule organizing center
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 400-660 of human FLII (NP_002009.1).
序列	LRLAGASPATVAAAAAAGSGPKDPMARKMRLRRRKDSAQDDQAKQVLKGMSDVAQEKNKKQESADARAPSGKVRR WDQGLEKPRLDYSEFFTEDVGLPGLTIWQIENFVPVLVEEAFHGKIFYEADCYIVLKTFLDDSGSLNWEIYYWIGGEATLD KKACSAIHAVNLRNYLGAECRTVREEMGDESEEFLLQVFDNDISYIEGGTASGFYTVEDTHYVTRMYRVYGGKKNIKLEPVPLK GTSLDPRFVFLDRGLDIYVWR

靶点信息

研究背景	This gene encodes a protein with a gelsolin-like actin binding domain and an N-terminal leucine-rich repeat-protein protein interaction domain. The protein is similar to a Drosophila protein involved in early embryogenesis and the structural organization of indirect flight muscle. The gene is located within the Smith-Magenis syndrome region on chromosome 17.
------	--

基因ID	2314
基因名	FLII
Swiss	Q13045
别名	FLII;FLI;FLIL;Fli1

产品验证



Western blot analysis of FLII expressed in A-549,293T,HeLa using FLII Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn