

# LIPH Rabbit pAb

货号: **AYP15452**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	50kDa
实测分子量	54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,DU145,293T,BxPC-3,A-549,Mouse testis,Mouse lung,Rat testis
细胞定位	Membrane,Peripheral membrane protein,Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 350-450 of human LIPH (NP_640341.1).
序列	KLRDKAGNTTESKINHEPTTFQKYHQVSLRLARFNQDLKVAASLMFSTGSLIGPRYKLRILRMKLRSLAHPERPOLCRYDLV LMENVETVFQPILCPQL

## 靶点信息

研究背景	This gene encodes a membrane-bound member of the mammalian triglyceride lipase family. It catalyzes the production of 2-acyl lysophosphatidic acid (LPA), which is a lipid mediator with diverse biological properties that include platelet aggregation, smooth muscle contraction, and stimulation of cell proliferation and motility.
基因ID	200879

基因名	LIPH
Swiss	Q8WWY8
别名	LIPH;AH;ARWH2;HYPT7;LAH2;LPDLR;PLA1B;mPA-PLA1;lipase H

## 产品验证



Western blot analysis of LIPH expressed in Mouse testis,Rat testis,HT-29 using LIPH Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)