

SWAP70 Rabbit pAb

货号: **AYP15458**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	68kDa
实测分子量	69kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,A-549,LO2
细胞定位	Cell membrane,Cell projection,Cytoplasm,Nucleus,lamellipodium
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 316-585 of human SWA P70 (NP_055870.2).
序列	HKEARQRRKELRKKQLAEQEELERQMKELQAANESKQQELEAVRKKLEEAASRAAEKKRLQTQVELQARFSTELEREKLI RQQMEEQVAQKSSELEQYLQRVRELEDMYKLEALEDERQARQDEETVRKLRLLLEEESKRAELEKWHLEQQQAIQT TEAEKQELNQRVLKEQALQEAMEQLELELERKQALEQYEEVKKKLEMATNKTQSWKDKVAHHEGLIRLIEPGSKNPHLIT NWWGPAAFTEAELEEREKNWKEKKTTE

靶点信息

研究背景	Phosphatidylinositol 3,4,5-trisphosphate-dependent guanine nucleotide exchange factor (GEF which, independently of RAS, transduces signals from tyrosine kinase receptors to RAC. It also mediates signaling of membrane ruffling. Regulates the actin cytoskeleton as an effector or adapter protein in response to agonist stimulated phosphatidylinositol (3,4-bisphosphate production and cell protrusion (By similarity.
------	---

基因ID	23075
基因名	SWAP70
Swiss	Q9UH65
别名	SWAP70;HSPC321;SWAP-70

产品验证



Western blot analysis of SWAP70 expressed in 293T, A-549, LO2 using SWAP70 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn