

PDXDC1 Rabbit pAb

货号: **AYP15459**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	47kDa/76kDa/83kDa/86kDa
实测分子量	87-106kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,A-549,HeLa,Mouse testis,Mouse brain,Rat heart
细胞定位	Golgi apparatus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 489-788 of human PDX DC1 (NP_055842.2).
序列	LCCTLQLREEFKQEVEATAGLLYVDDPNWSGIGVVRYEHANDDKSSLKSDPEGENIHAGLLKKLNELESDLTFKIGPEYKS MKSCLYVGMASDNVDAAELVETIAATAREIEENSRLLNMTEVVRKGIQEAQVELQKASEERLLEEGVLRQIPVVGSVLNWF SPVQALQKGRTFNLTAGSLESTEPIYVYKAQGAGVTLPTPSGSRTKQRLPGQKPKFRSLRGS DALSETSSVSHIEDLEKVE RLSSGPEQITLEASSTEGHPGAPSPQHTDQTEAFQKGVPHPEDDHSQVEGPESLR

靶点信息

研究背景	Enables cadherin binding activity. Predicted to be involved in carboxylic acid metabolic process. Located i n Golgi apparatus.
基因ID	23042

基因名	
Swiss	Q6P996
别名	PDXDC1;LP8165

产品验证



Western blot analysis of PDXDC1 expressed in Mouse testis,Rat heart,U-251MG using PDXDC1 Rabbit p Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 S.

实验步骤

访问官网浏览详情: www.ablybio.cn