

# APOBEC3C Rabbit pAb

货号: **AYP15482**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:50 - 1:100
理论分子量	22kDa
实测分子量	22kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	K-562
细胞定位	Cytoplasm, Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-190 of human APOBEC3C (NP_055323.2).
序列	MNPQIRNPMKAMYPGTFYFQFKNLWEANDRNETWLCFTVEGIKRRSVVSWKTGVFRNQVDSETHCHAERCFLSWFCDD ILSPNTKYQVTWYTSWSPDCAGEVAEFLARHSNVNLTIFTARLYYFYPCYQEGLRSLSQEGVAVEIMDYEDFKYCWEN FVYNDNEPFKPKWGLKTNFRLKRRRLRESLQ

## 靶点信息

研究背景	This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudo genes found in a cluster thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. It is thought that the proteins may be RNA editing enzymes and have roles in growth or cell cycle control.
------	--

基因ID	27350
基因名	APOBEC3C
Swiss	Q9NRW3
别名	APOBEC3C;A3C;APOBEC1L;ARDC2;ARDC4;ARP5;PBI;bK150C2.3

## 产品验证



Western blot analysis of APOBEC3C expressed in K-562 using APOBEC3C Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)