

— ABLYBIO, Help Your Research



DDI1 Rabbit pAb

货号: **AYP15485**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	44kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse testis
细胞定位	
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 97-396 of human DDI1 (NP_001001711.1).
------	--

靶点信息

研究背景	Probable aspartic protease (Probable. Seems to act as a proteasomal shuttle which links the proteasome and replication fork proteins like RTF2 (Probable. Required, with DDI2, for cellular survival following replication stress. Together or redundantly with DDI2, removes RTF2 from stalled forks to allow cell cycle progression after replication stress and maintains genome integrity.
基因ID	414301
基因名	DDI1
Swiss	Q8WTU0 (https://www.uniprot.org/uniprotkb/Q8WTU0/entry)
别名	DDI1,DDI1 Rabbit pAb

产品验证



Western blot analysis of DDI1 expressed in Mouse testis using DDI1 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)