

# UBQLNL Rabbit pAb

货号: **AYP15487**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	29kDa/52kDa
实测分子量	53kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse heart,Rat heart
细胞定位	cytosol
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 176-475 of human UBQLNL (NP_659490.4).
序列	LENPSIQRLLSNMEFMWQFISEHLDTQQLMQQNPEVSRLLLDNSEILLQTLELARNLAMIQEIMQIQPSQNLEYPLNPQPY LLETMPGGNNALGQNYADINDQMLNSMQDPFGGNPFTALLAGQVLEQVQSSPPPPPSQEQDQLTQHPATRVYINS SGGFSSNTSANDTLNKVNHTSKANTAMISTKGQSHICATRQPAWIPALPSIELTQQEQEYKDATVSLSSSRQTLKGDQL SDEQSSSQITGGMMQLLMNNPYLAAQIMLFTSMPQLSEQWRQQLPTFLQQTQISDLLSA

## 靶点信息

研究背景	Predicted to enable polyubiquitin modification-dependent protein binding activity. Predicted to be involved in ubiquitin-dependent protein catabolic process. Predicted to be active in cytosol.
基因ID	143630

基因名	UBQLNL
Swiss	Q8IYU4
别名	UBQLNL

## 产品验证



Western blot analysis of UBQLNL expressed in Mouse heart,Rat heart using UBQLNL Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)