

MIOX Rabbit pAb

货号: **AYP15491**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	24kDa/33kDa
实测分子量	33kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Rat kidney
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-285 of human MIOX (NP_060054.4).
序列	MKVTVGPDPSLVYRPDVDPEVAKDKASFRNYTSGPLLDREVFTTYKLMHHTHTVDFVRSKHAQFGGFSYKKMTVMEAVD LLDGLVDESDPDVDFPNSFHAFQTAEGIRKAHPDKDWFHLVGLLHDLGKVLALFGEPQWAVVGDTFVPGCRPQASVVF CDSTFQDNPDLDPRYSELGMYQPHCGLDRVLMSWGHDDEYMYQVMKFNKFSLPPEAFYMRHFHSFYPWHTGRDYQQ LCSQQDLAMLPPWVREFNKFELYTKCPDLPDVKLRPPYQGLIDKYCPGILSW

靶点信息

研究背景	Enables ferric iron binding activity and inositol oxygenase activity. Involved in inositol catabolic process. Predicted to be located in cytoplasm and inclusion body.
基因ID	55586

基因名	MIOX
Swiss	Q9UGB7
别名	MIOX;ALDRL6

产品验证



Western blot analysis of MIOX expressed in Mouse kidney,Rat kidney using MIOX Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn