

JAG2 Rabbit pAb

货号: **AYP15504**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	129kDa/133kDa
实测分子量	150kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse heart
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 24-200 of human JAG2 (NP_002217.3).
序列	ARPMGYFELQLSALRNVNGELLSGACCDGDGRTRTRAGGCGHDECPTYVVRVCLKEYQAKVTPTGPCSYGHGATPVLGGN SFYLPPAGAAGDRARARARAGGDQDPGLVIPFQFAWPRSFTLIVEAWDWDNDTTPNEELLIERVSHAGMINPEDRWKS LHFSGHVAHLELQIRVRCDE

靶点信息

研究背景	The Notch signaling pathway is an intercellular signaling mechanism that is essential for proper embryonic development. Members of the Notch gene family encode transmembrane receptors that are critical for various cell fate decisions. The protein encoded by this gene is one of several ligands that activate Notch and related receptors. Two transcript variants encoding different isoforms have been found for this gene.
------	---

基因ID	3714
基因名	JAG2
Swiss	Q9Y219
别名	JAG2;HJ2;SER2;jagged 2

产品验证



Western blot analysis of JAG2 expressed in Mouse heart using JAG2 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn