

RANKL Rabbit pAb

货号: **AYP15526**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	27kDa/30kDa/35kDa
实测分子量	31KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1,Mouse thymus,Rat thymus,Rat spleen
细胞定位	Cell membrane,Cytoplasm,Secreted,Single-pass type II membrane protein,Single-pass type II membrane protein
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 250-321 of human RANKL (NP_003692.1).
序列	PSSHTLMKGGSTKYWSGNSEHFHYSINVGGFKLRSGEEISIEVSNPSLLDPDQDATYFGAFKVRDID

靶点信息

研究背景	This gene encodes a member of the tumor necrosis factor (TNF) cytokine family which is a ligand for osteoprotegerin and functions as a key factor for osteoclast differentiation and activation. This protein was shown to be a dendritic cell survival factor and is involved in the regulation of T cell-dependent immune response. T cell activation was reported to induce expression of this gene and lead to an increase of osteoclastogenesis and bone loss. This protein was shown to activate antiapoptotic kinase AKT/PKB through a signaling complex involving SRC kinase and tumor necrosis factor receptor-associated factor (TRAF) 6, which indicated this protein may have a role in the regulation of cell apoptosis. Targeted disruption of the related gene in mice led to severe osteopetrosis and a lack of osteoclasts. The deficient mice exhibited defects in early differentiation of T and B lymphocytes, and failed to form lobulo-alveolar mammary structures during pregnancy. Two alternatively spliced transcript variants have been found.
基因ID	8600
基因名	TNFSF11
Swiss	O14788
别名	TNFSF11;CD254;ODF;OPGL;OPTB2;RANKL;TNLG6B;TRANCE;hRANKL2;sOdf

产品验证



Western blot analysis of RANKL expressed in Mouse thymus, Rat thymus, THP-1 using RANKL Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn