

# MAN1C1 Rabbit pAb

货号: **AYP15574**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	70kDa
实测分子量	68kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,U-87MG,HepG2,Mouse liver
细胞定位	Golgi apparatus membrane,Single-pass type II membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 44-240 of human MAN1 C1 (NP_065112.1).
序列	PHSSRLKRLFLAPRTQQPGLEVVAEIAGHAPAREQEPPNPAPAAPGEDDPSSWASPRRRKGLRRTRPTGPREEATAA RGNSIPASRPGDEGVPRFDNFNFRSRLRHPVLGTRADESQEPQSQVRAQREKIKEMMQFAWQSYKRYAMGKNELRPLT KDGYEGNMFGLSGATVIDSLDTLYLMELKEEFQEA

## 靶点信息

研究背景	Involved in the maturation of Asn-linked oligosaccharides. Trim alpha-1,2-linked mannose residues from Man(9GlcNAc(2 to produce first Man(8GlcNAc(2 then Man(6GlcNAc and a small amount of Man(5GlcNAc.
基因ID	57134

基因名	MAN1C1
Swiss	Q9NR34
别名	MAN1C1;HMIC;MAN1A3;MAN1C;pp6318

## 产品验证



Western blot analysis of MAN1C1 expressed in Mouse liver,A-549,U-87MG using MAN1C1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)