

— ABLYBIO, Help Your Research



Phospho-SQSTM1/p62-S349 Rabbit pAb

货号: **AYP15612**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:100 - 1:500
理论分子量	38kDa/47kDa
实测分子量	62KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,C6,NIH/3T3
细胞定位	Cytoplasm,Cytoplasmic vesicle,Endoplasmic reticulum,Late endosome,Lysosome,Nucleus,P-body,autophagosome
纯化	Affinity purification


抗原信息

抗原信息	A synthetic phosphorylated peptide around S349 of human SQSTM1/p62 (NP_003891.1).
------	---

靶点信息

研究背景	This gene encodes a multifunctional protein that binds ubiquitin and regulates activation of the nuclear factor kappa-B (NF- κ B) signaling pathway. The protein functions as a scaffolding/adaptor protein in concert with TNF receptor-associated factor 6 to mediate activation of NF- κ B in response to upstream signals. Alternatively spliced transcript variants encoding either the same or different isoforms have been identified for this gene. Mutations in this gene result in sporadic and familial Paget disease of bone.
基因ID	8878
基因名	SQSTM1
Swiss	Q13501 (https://www.uniprot.org/uniprotkb/Q13501/entry)
别名	SQSTM1,A170,DMRV,FTDALS3,NADGP,OSIL,PDB3,ZIP3,p60,p62,p62B,Phospho-SQSTM1/p62-S349 Rabbit pAb,EBI3-associated protein of 60 kDa,Phosphotyrosine-independent ligand for the Lck SH2 domain of 62 kDa,Ubiquitin-binding protein p62,ORCA

产品验证



Western blot analysis of Phospho-SQSTM1/p62-S349 expressed in HeLa, C6, NIH/3T3 using Phospho-SQSTM1/p62-S349 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)