

NTF4 Rabbit pAb

货号: **AYP15619**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC |
| 推荐浓度 | WB: 1:100 - 1:500 IHC: 1:100 - 1:200 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 22kDa |
| 实测分子量 | 22KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | MCF7,Mouse testis |
| 细胞定位 | Secreted |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 25-210 of human NTF4 (NP_006170.1). |
| 序列 | QPPPSTLPPFLAPEWDLSPRVVLSRGAPAGPPLLFLLEAGAFRESAGAPANRSRRGVSETAPASRRGELAVCDAVSGWVT DRRTAVDLRGREVEVLGEVPAAGGSPLRQYFFETRCKADNAEEGGPGAGGGGCRGVDRRHVWSECKAKQSYVRALTA DAQGRVGVWRWIRIDTACVCTLLSRTGRA |

靶点信息

| | |
|------|--|
| 研究背景 | This gene is a member of a family of neurotrophic factors, neurotrophins, that control survival and differentiation of mammalian neurons. The expression of this gene is ubiquitous and less influenced by environmental signals. While knock-outs of other neurotrophins including nerve growth factor, brain-derived neurotrophic factor, and neurotrophin 3 prove lethal during early postnatal development, NTF5-deficient mice only show minor cellular deficits and develop normally to adulthood. |
|------|--|

| | |
|-------|--|
| 基因ID | 4909 |
| 基因名 | NTF4 |
| Swiss | P34130 |
| 别名 | NTF4;GLC10;GLC10;NT-4;NT-4/5;NT-5;NT4;NT5;NTF5 |

产品验证



Western blot analysis of NTF4 expressed in MCF7, Mouse testis using NTF4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn